

**The Fermion Doubling Problem and Noncommutative Geometry**A. P. Balachandran\*, T.R.Govindarajan<sup>+</sup>, B. Ydri\* .*\*Physics Department, Syracuse University,  
Syracuse, N.Y., 13244-1130, U.S.A.**<sup>+</sup>Institute of Mathematical Sciences,  
Chennai 600 113, India***Abstract**

We propose a resolution for the fermion doubling problem in discrete field theories based on the fuzzy sphere and its Cartesian products.

The nonperturbative formulation of chiral gauge theories is a long standing programme in particle physics. It seems clear that one should regularise these theories with all symmetries intact. There are two major problems associated with conventional lattice approaches to this programme, both with roots in topological features: (1) The well-known Nielsen-Ninomiya theorem [1] states that if we want to maintain chiral symmetry, then under plausible assumptions, one cannot avoid the doubling of fermions in the usual lattice formulations . (2) It is not straightforward to understand features like anomalies with naive ultraviolet cut-off.

Recently a novel approach to discrete physics has been developed. It works with quantum fields on a “fuzzy space”  $\mathcal{M}_F$  obtained by treating the underlying manifold  $\mathcal{M}$  as a phase space and quantising it [2–9]. The earliest contributions to topological features of the emergent fuzzy physics came from Grosse, Klimčík and Prešnajder [4]. They dealt with monopoles and chiral anomaly for the fuzzy two-sphere  $S_F^2$  and took particular advantage of supersymmetry . Later Baez et al. [10] further elaborated on their monopole work and developed the fuzzy physics of  $\sigma$ – models and their solitons using cyclic cohomology [11, 12]. An attractive feature of this cohomological approach

is its ability to write analogues of continuum winding number formulae and derive a fuzzy Belavin-Polyakov bound [10].

In this paper, we propose a solution of the fermion doubling problem for  $\mathcal{M} = S^2$  using fuzzy physics. An alternative approach, also using fuzzy physics, can be found in [4] while further developments of our method with applications to instantons and chiral anomaly is reported in [13]. There have also been important developments [14] in the theory of chiral fermions and anomalies in the usual lattice formulations. They avoid fermion doubling by relaxing the requirement that chirality and Dirac operators anticommute. In contrast, fuzzy physics needs no such relaxation.

Quantisable adjoint orbits of compact semi-simple Lie groups seem amenable to the full fuzzy treatment and lead to manageable finite dimensional matrix models for quantum fields. There are two such manifolds in dimension four, namely  $S^2 \times S^2$  and  $\mathbb{CP}^2$ . Our methods readily extend to  $S^2 \times S^2$ . They are not anticipated to encounter obstructions for  $\mathbb{CP}^2$  as well. But we have not yet fully worked out its noncommutative geometry. The published work of Grosse and Strohmaier [15] on  $\mathbb{CP}^2$  gives their description of fuzzy 4d fermions.

A sphere  $S^2$  is a submanifold of  $\mathbb{R}^3$ :

$$S^2 = \langle \vec{n} \in \mathbb{R}^3 : \sum_{i=1}^3 n_i^2 = 1 \rangle. \quad (1)$$

If  $\hat{n}_i$  are the coordinate functions on  $S^2$ ,  $\hat{n}_i(\vec{n}) = n_i$ , then  $\hat{n}_i$  commute and the algebra  $\mathcal{A}$  of smooth functions they generate is commutative. In contrast, the operators  $x_i$  describing  $S_F^2$  are noncommutative:

$$[x_i, x_j] = \frac{i\epsilon_{ijk}x_k}{[l(l+1)]^{1/2}}, \quad \sum_{i=1}^3 x_i^2 = \mathbf{1}, \quad l \in \{\frac{1}{2}, 1, \frac{3}{2}, \dots\}. \quad (2)$$

The  $x_i$  commute and become  $\hat{n}_i$  in the limit  $l \rightarrow \infty$ . If  $L_i = [l(l+1)]^{1/2}x_i$ , then  $[L_i, L_j] = i\epsilon_{ijk}L_k$  and  $\sum L_i^2 = l(l+1)$ , so that  $L_i$  give the irreducible representation (IRR) of the  $SU(2)$  Lie algebra for angular momentum  $l$ .  $L_i$  or  $x_i$  generate the algebra  $A = M_{2l+1}$  of  $(2l+1) \times (2l+1)$  matrices.

Scalar wave functions on  $S^2$  come from elements of  $\mathcal{A}$ . In a similar way, elements of  $A$  assume the role of scalar wave functions on  $S_F^2$ . A scalar product on  $A$  is  $\langle \xi, \eta \rangle = \text{Tr} \xi^\dagger \eta$ .  $A$  acts on this Hilbert space by left- and right- multiplications giving rise to the left and right- regular representations

$A^{L,R}$  of  $A$ . For each  $a \in A$ , we thus have operators  $a^{L,R} \in A^{L,R}$  acting on  $\xi \in A$  according to  $a^L \xi = a\xi, a^R \xi = \xi a$ . [Note that  $a^L b^L = (ab)^L$  while  $a^R b^R = (ba)^R$ .] We assume by convention that elements of  $A^L$  are to be identified with fuzzy versions of functions on  $S^2$ . Note that there are two kinds of angular momentum operators  $L_i^L$  and  $-L_i^R$  for  $S_F^2$ . The orbital angular momentum operator, which should annihilate  $\mathbf{1}$ , is  $\mathcal{L}_i = L_i^L - L_i^R$ .  $\vec{\mathcal{L}}$  plays the role of the continuum  $-i(\vec{r} \times \vec{\nabla})$ .

The construction of the Dirac operator is of crucial importance for fuzzy physics. The following two Dirac operators on  $S^2$  have occurred in the fuzzy literature :

$$\mathcal{D}_1 = \vec{\sigma} \cdot [-i(\vec{r} \times \vec{\nabla})] + \mathbf{1}, \quad (3)$$

$$\mathcal{D}_2 = \epsilon_{ijk} \sigma_i \hat{n}_j \mathcal{J}_k, \quad (4)$$

where

$$\mathcal{J}_k = [-i(\vec{x} \times \vec{\nabla})]_k + \sigma_k/2 = \text{Total angular momentum operators} . \quad (5)$$

There is a common chirality operator  $\Gamma$  anticommuting with both:

$$\Gamma = \vec{\sigma} \cdot \hat{n} = \Gamma^\dagger, \quad \Gamma^2 = \mathbf{1}, \quad (6)$$

$$\Gamma \mathcal{D}_\alpha + \mathcal{D}_\alpha \Gamma = 0. \quad (7)$$

These Dirac operators *in the continuum* are unitarily equivalent,

$$\begin{aligned} \mathcal{D}_2 &= \exp(i\pi\Gamma/4) \mathcal{D}_1 \exp(-i\pi\Gamma/4) \\ &= i\Gamma \mathcal{D}_1, \end{aligned} \quad (8)$$

and have the spectrum

$$\text{Spectrum of } \mathcal{D}_\alpha = \{\pm(j+1/2) : j \in \{1/2, 3/2, \dots\}\}, \quad (9)$$

where  $j$  is total angular momentum (spectrum of  $\vec{\mathcal{J}}^2 = \{j(j+1)\}$ ).

Since  $|\mathcal{D}_\alpha|$  ( $\equiv$  positive square root of  $\mathcal{D}_\alpha^2$ ) for both  $\alpha$  share the same spectrum and eigenvectors by (9) and rotational invariance,  $|\mathcal{D}_1| = |\mathcal{D}_2|$ . Further being multiples of unity for each fixed  $j$ , they commute with the rotationally invariant  $\Gamma$ . As they are invertible too, we have the important identity

$$\Gamma = i \frac{\mathcal{D}_1}{|\mathcal{D}_1|} \frac{\mathcal{D}_2}{|\mathcal{D}_2|}. \quad (10)$$

The discrete version of  $\mathcal{D}_1$  is:

$$D_1 = \vec{\sigma} \cdot \vec{\mathcal{L}} + \mathbf{1}. \quad (11)$$

Its spectrum is

$$\begin{aligned} \text{Spectrum of } D_1 &= \left\{ \pm(j + \frac{1}{2}) : j \in \{\frac{1}{2}, \frac{3}{2}, \dots, 2l - \frac{1}{2}\} \right\} \\ &\cup \left\{ (j + \frac{1}{2}) : j = 2l + \frac{1}{2} \right\}. \end{aligned} \quad (12)$$

It is easy to derive (12) by writing

$$D_1 = \vec{J}^2 - \vec{\mathcal{L}}^2 - \left(\frac{\vec{\sigma}}{2}\right)^2 + \mathbf{1}, \quad (13)$$

$$\left(\frac{\vec{\sigma}}{2}\right)^2 = \frac{3}{4}\mathbf{1}, \quad (14)$$

$$\mathcal{L}_k + \frac{\sigma_k}{2} = J_k = \text{Total angular momentum operators.} \quad (15)$$

We let  $j(j+1)$  denote the eigenvalues of  $\vec{J}^2$ . Then for  $\vec{\mathcal{L}}^2 = k(k+1)$ ,  $k \in \{0, 1, \dots, 2l\}$ , if  $j = k + 1/2$  we get  $+(j + 1/2)$  as eigenvalue of  $D_1$ , while if  $j = k - 1/2$  we get  $-(j + 1/2)$ . The absence of  $-(2l + 1/2)$  in (12) is just because  $k$  cuts off at  $2l$ . (The same derivation works also for  $\mathcal{D}_1$ ).

The discrete version of  $\mathcal{D}_2$  is :

$$D_2 = -\epsilon_{ijk} \sigma_i x_j^L J_k = \epsilon_{ijk} \sigma_i x_j^L L_k^R. \quad (16)$$

$D_2$  is no longer unitarily equivalent to  $D_1$ , its spectrum being [7, 8]

$$\begin{aligned} \text{Spectrum of } D_2 &= \left\{ \pm(j + 1/2) \left[ 1 + \frac{1 - (j + 1/2)^2}{4l(l+1)} \right]^{1/2} : \right. \\ &\quad \left. j \in \{\frac{1}{2}, \frac{3}{2}, \dots, 2l + \frac{1}{2}\} \right\} \cup \{0 : j = 2l + \frac{1}{2}\}. \end{aligned} \quad (17)$$

The first operator has been used extensively by Grosse et al [4–6] while the second was first introduced by Watamuras [7, 8]. It is remarkable that the eigenvalues (12) coincide *exactly* with those of  $\mathcal{D}_\alpha$  upto  $j = (2l - 1/2)$ . In contrast  $D_2$  has zero modes when  $j = 2l + \frac{1}{2}$  and very small eigenvalues for large values of  $j$ , both being absent for  $\mathcal{D}_\alpha$ . So  $D_1$  is a better approximation to  $\mathcal{D}_\alpha$ .

But  $D_1$  as it stands admits no chirality operator anti-commuting with it. This is easy to see as its top eigenvalue does not have its negative in the spectrum. Instead  $D_2$  has the nice feature of admitting a chirality operator: the eigenvalue for top  $j$ , even though it has no pair, is exactly zero. So the best fuzzy Dirac operator has to combine the good properties of  $D_1$  and  $D_2$ . We suggest it to be  $D_1$  after projecting out its top  $j$  mode. We will show that it then admits a chirality with the correct continuum limit.

The chirality operator anticommuting with  $D_2$  and squaring to  $\mathbf{1}$  in the *entire* Hilbert space is

$$\gamma_2 = \gamma_2^\dagger = -\frac{\vec{\sigma} \cdot \vec{L}^R - 1/2}{l + 1/2}, \quad (18)$$

$$\gamma_2^2 = \mathbf{1}. \quad (19)$$

An interpretation of  $\gamma_2$  is that  $(1 \pm \gamma_2)/2$  are projectors to subspaces where  $(-\vec{L}^R + \vec{\sigma}/2)^2$  have values  $(l \pm \frac{1}{2})(l \pm \frac{1}{2} + 1)$  [10]. From (16) follows the identity

$$[D_1, \gamma_2] = -2i\lambda D_2; \lambda = \sqrt{1 - \frac{1}{(2l + 1)^2}}. \quad (20)$$

Now  $D_\alpha^2$  and  $|D_\alpha|$  ( $\equiv$  nonnegative square root of  $D_\alpha^2$ ) are multiples of identity for each fixed  $j$ , and  $\gamma_2$  commutes with  $\vec{J}$ . Hence they mutually commute :

$$[A, B] = 0 \text{ for } A, B = D_\alpha^2, |D_\alpha| \text{ or } \gamma_2. \quad (21)$$

Therefore from (20),

$$\{D_1, D_2\} = \frac{1}{2i\lambda} [D_1^2, \gamma_2] = 0. \quad (22)$$

In addition we can see that

$$[D_\alpha^2, D_\beta] = [|D_\alpha|, D_\beta] = 0. \quad (23)$$

If

$$\begin{aligned} \epsilon_\alpha &= \frac{D_\alpha}{|D_\alpha|} \quad \text{on the subspace } V \text{ with } j \leq 2l - 1/2, \\ &= 0 \quad \text{on the subspace } W \text{ with } j = 2l + 1/2. \end{aligned} \quad (24)$$

it follows that

$$e_1 = \epsilon_1, e_2 = \epsilon_2, e_3 = i\epsilon_1\epsilon_2 \quad (25)$$

generate a Clifford algebra on  $V$ . That is, if  $P$  is the orthogonal projector on  $V$ ,

$$\begin{aligned} P\xi &= \xi, \quad \xi \in V, \\ &= 0, \quad \xi \in W, \end{aligned} \quad (26)$$

then

$$\{e_\alpha, e_\beta\} = 2\delta_{\alpha\beta}P. \quad (27)$$

All this allows us to infer that  $\{e_3, D_1\} = 0$  so that it is a chirality operator for either  $D_1$  or its restriction  $PD_1P$  to  $V$ . In addition, *in view of (10, 25), it has the correct continuum limit as well* so that it is a good choice for chirality in that respect too.

A unitary transformation of  $e_3$  and  $D_1$  will not disturb their nice features. Such a transformation bringing  $e_3$  to  $\gamma_2$  on  $V$  is convenient. It can be constructed as follows.  $e_\alpha$  and  $\gamma_2$  being rotational scalars leave the two-dimensional subspaces in  $V$  with fixed values of  $\vec{J}^2$  and  $J_3$  invariant. On this subspace,  $e_\alpha$  and unity form a basis for linear operators, so  $\gamma_2$  is their linear combination. As  $e_1, e_3$  and  $\gamma_2$  anticommute with  $e_2$ , and all square to  $\mathbf{1}$ , in this subspace, we infer that  $\gamma_2$  is a transform by a unitary operator  $U = \exp(i\theta e_2/2)$  of  $e_3$  in each such subspace. And  $\theta$  can depend only on

$\vec{J}^2$  by rotational invariance. Thus we can replace  $PD_1P$  and  $e_3$  by the new Dirac and chirality operators

$$\begin{aligned} D &= e^{(i\theta(J^2)\epsilon_2)/2}(PD_1P)e^{(-i\theta(J^2)\epsilon_2)/2}, \\ \gamma &= P\gamma_2P = e^{(i\theta(J^2)\epsilon_2)/2}(i\epsilon_1\epsilon_2)e^{(-i\theta(J^2)\epsilon_2)/2} \\ &= \cos\theta(J^2)(i\epsilon_1\epsilon_2) + \sin\theta(J^2)\epsilon_1. \end{aligned} \quad (28)$$

The coefficients can be determined by taking traces with  $\epsilon_1$  and  $i\epsilon_1\epsilon_2$ .

We have established that chiral fermions can be defined on  $S_F^2$  with no fermion doubling, at least in the absence of fuzzy monopoles. We next extend this result to include them as well.

*Monopoles and Instantons:* In the continuum, monopoles and instantons are particular connection fields  $\omega$  on certain twisted bundles over the base manifold  $\mathcal{M}$ . On  $S^2$ , they are monopole bundles, on  $S^4$  or  $\mathbb{CP}^2$ , they can be  $SU(2)$  bundles.

In algebraic  $K$ -theory, it is well-known that these bundles are associated with projectors  $\mathcal{P}$  [11, 12, 16].  $\mathcal{P}$  is a matrix of some dimension  $M$  with  $\mathcal{P}_{ij} \in \mathcal{A} \equiv \mathcal{C}^\infty(\mathcal{M})$ ,  $\mathcal{P}^2 = \mathcal{P} = \mathcal{P}^\dagger$ . The physical meaning of  $\mathcal{P}$  is the following. Let  $\mathcal{A}^M = \mathcal{A} \otimes \mathbb{C}^M = \{\xi = (\xi_1, \xi_2 \dots \xi_M) : \xi_i \in \mathcal{A}\}$ . Then  $\mathcal{P}\mathcal{A}^M = \{\mathcal{P}\xi : \xi \in \mathcal{A}^M\}$  consists of smooth sections (or wave functions)  $\mathcal{P}\xi$  of a vector bundle over  $\mathcal{M}$ . For suitable choices of  $\mathcal{P}$ , we get monopole or instanton vector bundles. These projectors are known [16] and those for monopoles have been reproduced in [10].

The projectors  $p^{(\pm N)}$  for fuzzy monopoles of charge  $\pm N$  have also been explicitly found in [10]. They act on  $A^{2^N} = \{\xi \text{ with components } \xi_{b_1 \dots b_N} \in A, b_i \in \{1, 2\}\}$ . Let  $\vec{\tau}^{(i)}$  ( $i = 1, 2, \dots, N$ ) be commuting Pauli matrices.  $\vec{\tau}^{(i)}$  has the normal action on the index  $b_i$  and does not affect  $b_j$  ( $j \neq i$ ). Then  $\vec{K} = \vec{L}^L + \sum_i \vec{\tau}^{(i)}/2$  generates the  $SU(2)$  Lie algebra and  $p^{(N)}$  ( $p^{(-N)}$ ) is the projector to the maximum (minimum) angular momentum  $k_{\max} = l + N/2$  ( $k_{\min} = l - N/2$ ). [ $\vec{K}^2 p^{(N)} = k_{\max}(k_{\max} + 1)p^{(N)}$ ,  $\vec{K}^2 p^{(-N)} = k_{\min}(k_{\min} + 1)p^{(-N)}$ .] Fuzzy analogues of monopole wave functions are  $p^{(\pm N)}A^{2^N}$ . When spin is included, we must enhance  $p^{(\pm N)}A^{2^N}$  to  $p^{(\pm N)}A^{2^N} \otimes \mathbb{C}^2 = p^{(\pm N)}A^{2^{N+1}} = \{\xi \text{ with components } \xi_{b_1 \dots b_N, j} \in A : b_i, j \in \{1, 2\}\}$ .

The complications to be resolved now are caused by the need to project out a subspace of  $A^{2^{N+1}}$ . It is the analogue of the subspace projected out

by  $P$  for  $N = 0$ . In its absence, for example in the continuum, there is a canonical way due to Connes [11] for extending cyclic cohomology to twisted bundles.

In the  $N = 0$  sector, the projector  $P$  cuts out the subspace  $W$  of  $A^2$ . When we pass to  $p^{(\pm N)}A^{2N}$  and thence to  $p^{(\pm N)}A^{2N+1}$  by including spin, the subspace to be projected out is *not* determined by  $P$  if  $N \neq 0$ , as we shall see below. Rather, we can explain it as follows: Let  $\vec{J} = \vec{K} - \vec{L}^R + \vec{\sigma}/2$  be the “total angular momentum”. Calling  $\vec{J}$  by this name is appropriate as its components become (15) for  $N = 0$  and displays the known “spin-isospin mixing” [17] for  $N \neq 0$ . The maximum of  $\vec{J}^2$  on  $p^{(\pm N)}A^{2N+1}$  is  $J_{\max}(J_{\max} + 1)$ ,  $J_{\max} = (l \pm N/2) + l + 1/2 = 2l \pm N/2 + 1/2$ . [We assume that  $2l \geq (N-1)/2$ .] The vectors to be projected out are those with total angular momentum  $J_{\max}$ . If  $\mathcal{J}^{(\pm N)}$  are the corresponding projectors [with  $\mathcal{J}^{(0)} = P$ ], the twisted space we work with is  $\mathcal{J}^{(\pm N)}p^{(\pm N)}A^{2N+1}$ . Since  $p^{(\pm N)}$  commute with  $\vec{J}$  and hence with  $\mathcal{J}^{(\pm N)}$ ,  $Q^{(\pm N)} = \mathcal{J}^{(\pm N)}p^{(\pm N)}$  are also projectors.

There is no degeneracy for angular momentum  $J_{\max}$  in  $p^{(\pm N)}A^{2N+1}$ . That is because there is only one way to couple  $l \pm N/2, l$  and  $1/2$  to  $J_{\max}$ . The space  $(\mathbf{1} - \mathcal{J}^{(\pm N)})p^{(\pm N)}A^{2N+1}$  is thus of dimension  $2J_{\max} + 1$ . We want to get rid of this subspace.

The operators  $T = D$  or  $\gamma$  are zero on  $(\mathbf{1} - P)A^2$  where  $P$  cuts out states of angular momentum  $2l + 1/2$ . There is no degeneracy for this angular momentum in  $A^2$ .  $T$  and  $P$  extend canonically to  $A^2 \otimes \mathbb{C}^{2N} (\equiv A^{2N+1})$  as  $T \otimes \mathbf{1}$  and  $P \otimes \mathbf{1}$ . Let us call them once more as  $T$  and  $P$ .  $T$  and  $P$  commute with  $\vec{J}$  and hence with  $\mathcal{J}^{(\pm N)}$ . There is only one way to couple  $N$  “isospin”  $1/2$ ’s to  $(2l + 1/2)$  to get  $J_{\max}$  so that  $(\mathbf{1} - \mathcal{J}^{(\pm N)})(\mathbf{1} - P)A^{2N+1}$  is also of dimension  $2J_{\max} + 1$ . And  $T$  is zero on this subspace.

The projectors  $(\mathbf{1} - \mathcal{J}^{(\pm N)})p^{(\pm N)}$  and  $(\mathbf{1} - \mathcal{J}^{(\pm N)})(\mathbf{1} - P)$  being of the same rank, there exists a unitary operator  $U$  on  $A^{2N+1}$  transforming one to the other:

$$(\mathbf{1} - \mathcal{J}^{(\pm N)})p^{(\pm N)} = U(\mathbf{1} - \mathcal{J}^{(\pm N)})(\mathbf{1} - P)U^{-1}. \quad (29)$$

If we transport  $T$  by  $U$ ,

$$T' = UTU^{-1}, \quad (30)$$

then  $T' = D', F'$  or  $\gamma'$  vanishes on  $(\mathbf{1} - \mathcal{J}^{(\pm N)})p^{(\pm N)}A^{2N+1}$ . On its orthogonal complement  $[\mathbf{1} - (\mathbf{1} - \mathcal{J}^{(\pm N)})p^{(\pm N)}]A^{2N+1}$ , invariant under  $T'$ ,  $\gamma'$  squares to



unity and anticommutes with  $D'$  just as we want. What replaces  $P$  now is not  $\mathcal{J}^{(\pm N)}$ , but rather

$$P^{(\pm N)} = [\mathbf{1} - (\mathbf{1} - \mathcal{J}^{(\pm N)})p^{(\pm N)}], \quad (31)$$

$$P^{(0)} = P. \quad (32)$$

As  $l \rightarrow \infty$ ,  $J_{\max}$  becomes dominated by  $2l$  and so we have the freedom to let  $U$  approach  $\mathbf{1}$ . That is, no  $U$  is needed in the continuum limit.

Total angular momentum  $2l + 1/2 + N/2$  has no multiplicity in  $A^{2^{N+1}}$ . As both  $(\mathbf{1} - \mathcal{J}^{(N)})(\mathbf{1} - P)A^{2^{N+1}}$  and  $(\mathbf{1} - \mathcal{J}^{(N)})p^{(N)}A^{2^{N+1}}$  have this angular momentum, we have that

$$(\mathbf{1} - \mathcal{J}^{(N)})(\mathbf{1} - P)A^{2^{N+1}} = (\mathbf{1} - \mathcal{J}^{(N)})p^{(N)}A^{2^{N+1}}. \quad (33)$$

So we choose

$$U = \mathbf{1} \quad \text{on} \quad (\mathbf{1} - \mathcal{J}^{(N)})(\mathbf{1} - P)A^{2^{N+1}}. \quad (34)$$

Next in accordance with (29), we set

$$U(\mathbf{1} - \mathcal{J}^{(-N)})(\mathbf{1} - P)A^{2^{N+1}} = (\mathbf{1} - \mathcal{J}^{(-N)})p^{(-N)}A^{2^{N+1}}. \quad (35)$$

We also demand that

$$[U, J_i] = 0. \quad (36)$$

That fixes  $U$  upto a phase on the subspace  $(\mathbf{1} - \mathcal{J}^{(-N)})(\mathbf{1} - P)A^{2^{N+1}}$ .

We saw in [10] that  $(1 \pm \gamma)/2$  are projectors for combining  $-\vec{L}^R$  and  $\vec{\sigma}/2$  to give angular momenta  $l \pm 1/2$ . So  $\gamma = +1$  on all the subspaces  $(\mathbf{1} - \mathcal{J}^{(\pm N)})(\mathbf{1} - P)A^{2^{N+1}}$ ,  $(\mathbf{1} - \mathcal{J}^{(\pm N)})p^{(\pm N)}A^{2^{N+1}}$ . Also,  $(\frac{\sum \vec{\tau}^{(i)}}{2})^2$  is  $\frac{N}{2}(\frac{N}{2} + 1)$  on these subspaces. It follows that (34), (35) and (36) are compatible with a  $U$  commuting with  $J_i$ ,  $\gamma$ ,  $(-\vec{L}^R + \vec{\sigma}/2)^2$  and  $(\frac{\sum \vec{\tau}^{(i)}}{2})^2$ . We now outline an extension of  $U$  to all of  $A^{2^{N+1}}$  consistently with rotational invariance (36) and

$$\left[ U, \left( -\vec{L}^R + \frac{\vec{\sigma}}{2} \right)^2 \right] = \left[ U, \left( \frac{\sum \vec{\tau}^{(i)}}{2} \right)^2 \right] = 0. \quad (37)$$

An important consequence is that

$$\gamma' = \gamma \quad (38)$$

so that

$$[\gamma', p^{(\pm N)}] = [\gamma', J_i] = [\gamma', \mathcal{J}^{(\pm N)}] = 0. \quad (39)$$

One way to specify  $U$  more fully is as follows. Let

$$A^{2^{N+1}} = X \oplus X^\perp = X' \oplus X'^\perp \quad (40)$$

be orthogonal decompositions where

$$X = (\mathbf{1} - \mathcal{J}^{(N)})(\mathbf{1} - P)A^{2^{N+1}} \oplus (\mathbf{1} - \mathcal{J}^{(-N)})(\mathbf{1} - P)A^{2^{N+1}}, \quad (41)$$

$$X' = UX = (\mathbf{1} - \mathcal{J}^{(N)})p^{(N)}A^{2^{N+1}} \oplus (\mathbf{1} - \mathcal{J}^{(-N)})p^{(-N)}A^{2^{N+1}}. \quad (42)$$

Both  $X$  and  $X'$  are invariant under the self-adjoint operators

$$J_i, \left(-\vec{L}^R + \frac{\vec{\sigma}}{2}\right)^2 \quad \text{and} \quad \left(\sum \frac{\vec{\tau}^{(i)}}{2}\right)^2.$$

Therefore, the same is the case with  $X^\perp$  and  $X'^\perp$ . We can extend  $U$  to a map  $X^\perp \rightarrow X'^\perp$  which commutes with the above operators. There would still be uncertainties about choosing  $U$  requiring further conventions for elimination.

Although  $T'$  are operators on  $P^{(\pm N)}A^{2^{N+1}}$ , that is not the space of sections for the twisted bundles. The latter is, rather,

$$Q^{(\pm N)}A^{2^{N+1}} = p^{(\pm N)}P^{(\pm N)}A^{2^{N+1}} = p^{(\pm N)}\mathcal{J}^{(\pm N)}A^{2^{N+1}}. \quad (43)$$

It is not an invariant subspace for  $D'$  unless  $D'$  is projected, or corrected by connections as explained in [13]. However, chirality  $\gamma'$  is well-defined on twisted sections because of (39).

We now permanently rename  $T'$ ,  $P^{(\pm N)}$  and  $Q^{(\pm N)}$  as follows:

$$\begin{aligned} D', \gamma' &\rightarrow D, \gamma, \quad p^{(\pm N)} \rightarrow p, \\ P^{(\pm N)} &\rightarrow P, \quad Q^{(\pm N)} \rightarrow Q. \end{aligned} \quad (44)$$

$\gamma'$  in any case is  $\gamma$ .

This completes our construction of Dirac operator and chirality on  $S_F^2$ . We may remark here that the algebra for the space  $S_F^2 \times S_F^2$  is  $A \otimes_{\mathbb{C}} A$  while its Dirac and chirality operators are  $D \otimes \mathbf{1} + \gamma \otimes D$  and  $\gamma \otimes \gamma$ .

### Acknowledgments

S.Vaidya, Xavier Martin, Denjoe O'Connor and Peter Prešnajder offered us many good suggestions during this work while Apoorva Patel told us about [14]. We thank them for their help. The work of APB was supported in part by the DOE under contract number DE-FG02-85ER40231.

## References

- [1] N. B. Nielsen and M. Ninomiya, *Phys.Lett.*, B105: 219, 1981; *Nucl.Phys.*, B185: 20, 1981.
- [2] J. Madore. *An Introduction to Noncommutative Differential Geometry and its Applications*. Cambridge University Press, Cambridge, 1995; [gr-qc/9906059](#).
- [3] H. Grosse and P. Prešnajder. *Lett.Math. Phys.*, 33:171–182, 1995. and references therein.
- [4] H. Grosse, C. Klimčík, and P. Prešnajder. *Commun.Math.Phys.*, **178**:507–526, 1996; **185**:155–175, 1997; H. Grosse and P. Prešnajder. *Lett.Math.Phys.* **46**:61–69, 1998 and ESI preprint, 1999.
- [5] H. Grosse, C. Klimčík, and P. Prešnajder. *Commun.Math.Phys.*, 180:429–438, 1996. [hep-th/9602115](#).
- [6] H. Grosse, C. Klimčík, and P. Prešnajder. In *Les Houches Summer School on Theoretical Physics*, 1995. [hep-th/9603071](#). See citations in [3–6] for further references.
- [7] U. Carow-Watamura and S. Watamura. [hep-th/9605003](#). *Commun.Math.Phys.*,183:365–382, 1997.
- [8] U. Carow-Watamura and S. Watamura. [hep-th/9801195](#).
- [9] J. Frohlich, O. Grandjean, and A. Recknagel. *Commun. Math. Phys.*, 203:119–184,, 1999. [math-phy/9807006](#) and references therein for fuzzy group manifolds.
- [10] S. Baez, A. P. Balachandran, S. Vaidya and B. Ydri. [hep-th/9811169](#) and *Comm.Math.Phys* (in press).
- [11] A. Connes. *Noncommutative Geometry*. Academic Press, London, 1994; G. Landi. *An Introduction to Noncommutative Spaces And Their Geometries*. Springer-Verlag , Berlin , 1997 . [hep-th/9701078](#).
- [12] R. Coquereaux. *J. Geo. Phys.*, 6:425–490, 1989.

- [13] A. P. Balachandran, S. Vaidya. *SU-4210-701*, *TIFR/TH/99-32* and *hep-th/9910129*.
- [14] M. Luscher. *Phys. Lett.* **B428** (1998) 342 and *hep-lat/9802011* ; *hep-lat/9904009* and references therein .
- [15] H. Grosse and A. Strohmaier. *Lett.Math.Phys.* **48**:163–179,1999. *hep-th/9902138*.
- [16] J. A. Mignaco, C. Sigaud, A. R. da Silva and F. J. Vanhecke. *Rev.Math.Physics*, **9** (1997) 689 and *hep-th/9611058*; G. Landi. *math-phy/9812004*; *math-phy/9905014*.
- [17] R. Jackiw and C. Rebbi. *Phys.Rev.Lett.* **36**:1116-1119, 1976 ; P. Hasenfratz and G. 't Hooft. *Phys.Rev.Lett.* **36**:1119-1122, 1976.